

Exercise: Wannier Functions

We will construct maximally localized Wannier functions for SrVO_3 (or a material of your choice) using *wien2wannier* and *wannier90*.

NB: there is no *w2web* module for this. You have to use the command line.

NB2: Steps 1–8 will guide you through the complete “Wannierization” process. Everything after that is “optional”.

More information:

http://www.wien2k.at/reg_user/unsupported/wien2wannier/
→ user guide

<http://wannier.org>

Easy as 1-2-3 ...

- 1 **standard Wien2k** run
 - spacegroup 221 $Pm - 3m$, $a = 7.2613$ Bohr
 - Sr (0, 0, 0), V ($1/2, 1/2, 1/2$), O ($0, 1/2, 1/2$)
 - obtain **band structure**
- 2 identify **target bands** (“low-energy model”)
 - minimal (d -only): $V-t_{2g}$ around E_F
 - or (your choice) dp -model: $V-t_{2g} + O-p$
- 3 **prepare** a directory for wien2wannier
 - decide on a k -mesh (e.g. $4 \times 4 \times 4$)
 - `prepare_w2wdir.sh case w2wdir`
creates `w2wdir` (use e.g. k444) and copies some files
 - `cd w2wdir`

... 4-5-6-7-8

- 4 write input files: `init_w2w`
 - follow `instructions`
 - do `not shift k-mesh`
 - use “common sense” ...
- 5 eigenvectors on full k-mesh: `x lapw1`
- 6 compute overlap matrix: `w2w case`
- 7 set $E_F \leftarrow 0$: `shift_energy case`
- 8 optimize $U(\mathbf{k})$: `wannier90.x case`

...and you're done!

Consistency Checks and Analysis

- Wannier function spread $\langle \Delta r^2 \rangle$
 - look for “Final State” in `case.wout`
 - should be $\sim 1.6 \text{ \AA}^2$
- compare **bandstructure** Wien2k vs. Wannier interpolation
 - in gnuplot:
plot 'case.spaghetti_ene' using (\$4/0.35):5,
 '**case_band.dat**' with lines
- look at the **hopping** matrix elements
(\leftrightarrow tight-binding Hamiltonian $H(\mathbf{R})$, `case_hr.dat`)
 - Do you see the proper **symmetries**?
 - Do the hoppings **decay** quickly enough?
(Exponential localization!)

If You Still Have Time

- compare *d-only* and *dp-models*
 - How do the “V-d” WF change (hoppings, spread)?
- Check **k-mesh convergence**. When do you see convergence of
 - the bandstructure?
 - $H(\mathbf{R})$?
 - the spreads?
- **plot** the Wannier functions (→ user guide, or ask help)
 - ① `write_wplotin case` ① `prepare_plots.sh case`
 - ② `write_wplotdef case` ② `xsfAll.sh case`
 - ③ `wplot case` ③ `xcrysden -xsf case_i.xsf.gz`
 - ...